

CHECKLIST OF LANCASHIRE AND CHESHIRE SPIDERS

The following checklist of Lancashire (VC 59 and 60) and Cheshire (VC 58) spiders has been compiled from a variety of sources especially Falconer (1930), Bristowe (1939) Mackie (1960 et. seq.), nomenclature follows Roberts (1987). The total species count stands at 320 for Lancashire and 343 for Cheshire. A number of species listed by Mackie in his Lancashire & Cheshire Fauna Society reports are from VC 69 and these have been omitted from this checklist. These species are *Dolomedes fimbriatus* (Clerck), *Clubiona coeruleascens* L.Koch, *Marolepidus casemeir*, and *Theridion bellicosum* Simon. Roberts (1987) has shown that several spiders holding specific status are in fact varieties of other species. These are *Pelecopsis locketi* Cooke, = *P. nemoralis* (Blackwall), *Pelecopsis mediocris* Kulczynski = *P. nemoralis* (Blackwall), *Dicymbium brevisetosum* Locket = *D. nigrum* (Blackwall), *Oedothorax tuberosus* (Blackwall) = *O. gibbosus* (Blackwall) and *Troxochrus cirrifrons* (O.P.- Cambridge) = *T. scabriculus* (Westring) respectively.

Doubtful records:

Mackie (1975) throws doubt on the validity of *Liocranum rupicola* (Walckenaer) from Poynton, Cheshire 1903 and *Mangora acalypha* (Walckenaer) from Chat Moss, Lancashire in 1900 after examining the H.W.Freston Collection at Manchester Museum. He also expressed doubts (pers. comm.) about *Marpissa pomatia* (Clerck) recorded at Southport in 1855, in view of its known distribution in Britain. *Dysdera erythrina* (Walckenaer) which has not been recorded since the Manchester record of 1842 may also be doubtful.

Probable Introductions:

Pholcus phalangioides (Fuesslin), *Physocyclus simoni* (Berland), *Scytodes thoracica* Latreille, and *Micaria subopaca* Westring, are included in the list even though it is very probable that all of them have been accidentally imported into the area.

Unpublished new records for Lancashire:

(LM) = Liverpool Museum specimen.

Scytodes thoracica Latreille Liverpool (SJ 35-89-) C.Felton, 04.10. 1991 (LM); *Pirata uliginosus* (Thorell) Rixton Moss (SJ 66-90-), C.R.Billups, 09.08.1902 (two specimens in Warrington Museum ante-date the previous first recorded specimen at Kirkby Moss of

07.1969); *Enoplognatha latimana* Hippa & Oksala, Freshfield (SD 28-09-), S. Dobson, 08.1984; (LM); *Larinoides sclopetaarius* (Clerck), Fiddlers Ferry (SJ 56-86-), J. Wright, 1983; *Atea sturmi* (Hahn), Ainsdale (SD 28-10-), C. Felton, 08.85; *Walckenaeria capito* (Westring), Hale Duck-decoy (SJ 47-82-), C. Felton, 08.11.1976 (LM); *Walckenaeria incisa* (O.P.- Cambridge), Kirkby (SJ 43-98-), C. Felton, 05.12.1983 (LM); *erigone vagans* Audouin, Seaforth (SJ 31-96-), C. Felton, 19.09.1982 (LM); *Baryphyma trifrons* (O.P.- Cambridge), Alt Estuary (SD 29-03-), C. Felton, 03.1983 (LM); *Milleriana inerrans* (O.P.- Cambridge), Hightown (SD 29-03-), C. Felton, 31.05.1984 (LM); *Porrhomma pallidum* Jackson, Rainford Delf (SJ 51-99-), J. Wright, 30.04.89; *Saaristoa firma* (O.P.- Cambridge), Knowsley, (SJ 42-95-), C. Felton, 20.05.1971 (LM).

Unpublished new records for Cheshire:

Pirata tenuitarsus Simon, Little Budworth (SJ 58-65-), S. Dobson, 1982; *Enoplognatha latimana* Hippa & Oksala, Wallasey (SJ 27-92-), C. Felton, 07.08.1989 (LM); *Mecopisthes peusi* Wunderlich, Wallasey (SJ 27-92-), C. Felton, 1982 (LM); *Erigone vagans* Audouin, Hilbre (SJ 18-87-), C. Felton, 08.08.1982 (LM); *Agyneta olivacea* (Emerton), Flaxmere (SJ 55-72-), C. Felton, 08.1991 (LM).

Abbreviations: Species not recorded since Falconer (1930) *

Post 1960 records +

Doubtful records d

Introduced species x

No known record -

Lancashire L

Cheshire C

ATYPIDAE

Atypus affinis Eichwald L+ C+

AMAUROBIIDAE

Amaurobius fennestralis (Stroem) L+ C+

Amaurobius similis (Blackwall) L+ C+

Amaurobius ferox (Walckenaer) L+ C+

DICTYNIDAE

<i>Dictyna arundinacea</i> (Linnaeus)	L+	C+
<i>Dictyna uncinata</i> Thorell	L+	C+
<i>Dictyna latens</i> (Fabricius)	L+	C+
<i>Lathys humiliis</i> (Blackwall)	-	C*
<i>Argenna subnigra</i> (O.P.- Cambridge)	-	C*
<i>Argenna patula</i> (Simon)	L+	C

OONOPIDAE

<i>Oonops pulcher</i> Templeton	L+	C+
<i>Oonops domesticus</i> de Dalmas	L+	C+

SCYTODIDAE

<i>Scytodes thoracica</i> Latreille	L+	-
-------------------------------------	----	---

DYSDERIDAE

<i>Dysdera erythrina</i> (Walckenaer)	L*	-
<i>Dysdera crocata</i> C.L. Koch	L+	C+
<i>Harpactea hombergi</i> (Scopoli)	L+	C+

SEGESTRIIDAE

<i>Segestria senoculata</i> (Linnaeus)	L+	C+
--	----	----

PHOLCIDAE

<i>Pholcus phalangioides</i> (Fuesslin)	L+	C
<i>Psilochorus simoni</i> (Berland)	-	C*x

GNAPHOSIDAE

<i>Drassodes lapidosus</i> (Walckenaer)	L+	C+
<i>Drassodes cupreus</i> (Blackwall)	L+	C+
<i>Drassodes pubescens</i> (Thorell)	L*	-
<i>Haplodrassus signifer</i> (C.L. Koch)	L+	C+
<i>Scotophaeus blackwalli</i> (Thorell)	L+	C+
<i>Zelotes electus</i> (C.L. Koch)	L+	-
<i>Zelotes latreillei</i> (Simon)	L+	C+
<i>Zelotes apricorum</i> (L. Koch)	L+	-
<i>Zelotes pusillus</i> (C.L.Koch)	L*	-
<i>Gnaphosa leporina</i> (L. Koch)	-	C+
<i>Micaria pulicaria</i> (Sundevall)	L+	C+
<i>Micaria subopaca</i> Westring	L+	-

CLUBIONIDAE

<i>Clubiona corticalis</i> (Walckenaer)	L+	C+
<i>Clubiona reclusa</i> O.P.- Cambridge	L+	C+

<i>Clubiona stagnatilis</i> Kulczynski	L+	C+
<i>Clubiona norvegica</i> Strand	-	C+
<i>Clubiona pallidula</i> (Clerck)	L+	C+
<i>Clubiona phragmitis</i> C.L. Koch	L+	C+
<i>Clubiona terrestris</i> Westring	L+	C+
<i>Clubiona neglecta</i> O.P.- Cambridge	L+	C+
<i>Clubiona lutescens</i> Westring	L+	C+
<i>Clubiona compta</i> C.L. Koch	L+	C+
<i>Clubiona brevipes</i> Blackwall	L+	C+
<i>Clubiona trivialis</i> C.L. Koch	L+	C+
<i>Clubiona diversa</i> O.P.- Cambridge	L+	C+
<i>Clubiona subtilis</i> L. Koch	L*	-
<i>Cheiracanthium erraticum</i> (Walckenaer)	L+	C+
<i>Cheiracanthium virescens</i> (Sundevall)	L+	C+
LIOCRANIDAE		
<i>Agroeca brunnea</i> (Blackwall)	L+	C+
<i>Agroeca proxima</i> (O.P.- Cambridge)	L+	C+
<i>Scotina celans</i> (Blackwall)	L+	-
<i>Scotina gracilipes</i> (Blackwall)	L+	C+
<i>Liocranum rupicola</i> (Walckenaer)	-	C*d
<i>Phrurolithus festivus</i> (C.L. Koch)	L*	C+
ZORIDAE		
<i>Zora spinimana</i> (Sundevall)	L+	C+
ANYPHAENIDAE		
<i>Anyphaena accentuata</i> (Walckenaer)	L+	-
EUSPARASSIDAE		
<i>Micrommata virescens</i> (Clerck)	L*	-
THOMISIDAE		
<i>Xysticus cristatus</i> (Clerck)	L+	C+
<i>Xysticus audax</i> (Schrank)	L	C+
<i>Xysticus kochi</i> Thorell	L	C
<i>Xysticus erraticus</i> (Blackwall)	L	C+
<i>Xysticus ulmi</i> (Hahn)	L	C
<i>Xysticus sabulosus</i> (Hahn)	L+	C+
<i>Oxyptila praticola</i> (C.L. Koch)	L	C
<i>Oxyptila trux</i> (Blackwall)	L+	C+
<i>Oxyptila atomaria</i> (Panzer)	L+	C+

Oxyptila brevipes (Hahn)	-	C+
PHILODROMIDAE		
Philodromus dispar Walckenaer	L+	C+
Philodromus aureolus (Clerck)	L+	C+
Philodromus cespitum (Walckenaer)	L+	C+
Philodromus fallax Sundevall	L+	-
Philodromus histrio (Latreille)	-	C+
Thanatus striatus C.L. Koch	L*	-
Tibellus maritimus (Menge)	L+	C+
Tibellus oblongus (Walckenaer)	L+	C+
SALTICIDAE		
Salticus scenicus (Clerck)	L+	C+
Salticus cingulatus (Panzer)	L+	C+
Heliophanus cupreus (Walckenaer)	L+	C+
Heliophanus flavipes (Hahn)	L+	C+
Marpissa muscosa (Clerck)	L*d	-
Marpissa radiata (Grube)	L*	-
Bianor aurocinctus (Ohlert)	L*	-
Neon reticulatus (Blackwall)	L+	C+
Euophrys frontalis (Walckenaer)	L+	C+
Euophrys herbigrada (Simon)	L+	-
Euophrys erratica (Walckenaer)	L+	C
Euophrys lanigera (Simon)	L+	C+
Sitticus pubescens (Fabricius)	L+	C+
Sitticus floricola (C.L. Koch)	-	C+
Attulus saltator (Simon)	L+	C+
Evarcha falcata (Clerck)	L+	C+
LYCOSIDAE		
Pardosa agricola (Thorell)	L+	C*
Pardosa purbeckensis (F.O.P.- Cambridge)	L+	C+
Pardosa monticola (Clerck)	L+	C+
Pardosa palustris (Linnaeus)	L+	C+
Pardosa prativaga (L. Koch)	L+	C+
Pardosa amentata (Clerck)	L+	C+
Pardosa nigriceps (Thorell)	L+	C+
Pardosa lugubris (Walckenaer)	L+	C+
Pardosa hortensis (Thorell)	L	-

<i>Xerolycosa minata</i> (C.L. Koch)	L+	C+
<i>Alopecosa pulverulenta</i> (Clerck)	L+	C+
<i>Alopecosa accentuata</i> (Latreille)	L+	-
<i>Trochosa ruricola</i> (Degeer)	L+	C+
<i>Trochosa terricola</i> Thorell	L+	C+
<i>Trochosa spinipalpis</i> (F.O.P.- Cambridge)	L+	C+
<i>Arctosa perita</i> (Latreille)	L+	C+
<i>Arctosa leopardus</i> (Sundevall)	L+	C+
<i>Arctosa cinerea</i> (Fabricius)	L+	-
<i>Pirata piscatorius</i> (Clerck)	L+	C+
<i>Pirata piraticus</i> (Clerck)	L+	C+
<i>Pirata hygrophilus</i> Thorell	L+	C+
<i>Pirata uliginosus</i> (Thorell)	L+	C+
<i>Pirata latitans</i> (Blackwall)	L+	C+
<i>Pirata tenuitarsus</i> Simon	-	C+
PISAURIDAE		
<i>Pisaura mirabilis</i> (Clerck)	L+	C+
ARGYRONETIDAE		
<i>Argyroneta aquatica</i> (Clerck)	L+	C+
AGELENIDAE		
<i>Agelena labyrinthica</i> (Clerck)	L+	C+
<i>Tetrix denticulata</i> (Olivier)	L+	C+
<i>Tegenaria gigantea</i> Chamberlin & Ivie	L+	C+
<i>Tegenaria sacra</i> Blackwall	L+	C+
<i>Tegenaria atrica</i> C.L. Koch	L+	C+
<i>Tegenaria agrestis</i> (Walckenaer)	L+	C+
<i>Tegenaria domestica</i> (Clerck)	L+	C+
<i>Tegenaria silvestris</i> L. Koch	L+	C+
<i>Cocloetes atropos</i> (Walckenaer)	L+	C+
<i>Cryphoeca silvicola</i> (C.L. Koch)	L+	C+
HAHNIIDAE		
<i>Antistea elegans</i> (Blackwall)	L+	C+
<i>Hahnia montana</i> (Blackwall)	L+	C+
<i>Hahnia nava</i> (Blackwall)	L+	C+
<i>Hahnia helveola</i> Simon	L+	C+
<i>Hahnia pusilla</i> C.L. Koch	L+	C+

MIMETIDAE

<i>Ero cambridgei</i> Kulczynski	L+	C+
<i>Ero furcata</i> (Villers)	L+	C+

THERIDIIDAE

<i>Episinus angulatus</i> (Blackwall)	L+	C+
<i>Euryopis flavomaculata</i> (C.L. Koch)	L*	C+
<i>Crustulina guttata</i> (Wider)	-	C+
<i>Steatoda phalerata</i> (Panzer)	-	C+
<i>Steatoda bipunctata</i> (Linnaeus)	L+	C+
<i>Steatoda grossa</i> (C.L. Koch)	-	C+
<i>Anelosimus vittatus</i> (C.L. Koch)	L+	C+
<i>Achaearanea tepidariorum</i> (C.L. Koch)	L*	C*
<i>Theridion sisyphium</i> (Clerck)	L+	C+
<i>Theridion impressum</i> L. Koch	-	C+
<i>Theridion pictum</i> (Walckenaer)	L+	C+
<i>Theridion varians</i> Hahn	L+	C+
<i>Theridion simile</i> C.L. Koch	-	C+
<i>Theridion melanurum</i> Hahn	L+	C+
<i>Theridion mystaceum</i> L. Koch	L+	C+
<i>Theridion blackwalli</i> O.P.- Cambridge	L*	C*
<i>Theridion tinctum</i> (Walckenaer)	-	C
<i>Theridion bimaculatum</i> (Linnaeus)	L+	C+
<i>Theridion pallens</i> Blackwall	L+	C+
<i>Enoplognatha ovata</i> (Clerck)	L+	C+
<i>Enoplognatha latimana</i> Hippa & Oksala	L+	C+
<i>Enoplognatha thoracica</i> (Hahn)	L+	C+
<i>Robertus lividus</i> (Blackwall)	L+	C+
<i>Robertus arundineti</i> (O.P.- Cambridge)	-	C+
<i>Robertus neglectus</i> (O.P.- Cambridge)	L*	C+
<i>Pholcomma gibbum</i> (Westring)	L+	C+
<i>Theonoe minutissima</i> (O.P.- Cambridge)	-	C+

NESTICIDAE

<i>Nesticus cellularis</i> (Clerck)	L+	C+
-------------------------------------	----	----

TETRAGNATIDAE

<i>Tetragnatha extensa</i> (Linnaeus)	L+	C+
<i>Tetragnatha montana</i> Simon	L+	C+
<i>Tetragnatha obtusa</i> C.L. Koch	L+	C+

<i>Tetragnatha nigrita</i> Lendl	-	C+
<i>Tetragnatha striata</i> L. Koch	-	C*
<i>Pachygynatha clercki</i> Sundevall	L+	C+
<i>Pachygynatha degeeri</i> Sundevall	L+	C+
<i>Pachygynatha listeri</i> Sundevall	L	C
METIDAE		
<i>Metellina segmentata</i> (Clerck)	L+	C+
<i>Metellina mengci</i> (Blackwall)	L+	C+
<i>Metellina merianae</i> (Scopoli)	L+	C+
<i>Meta menardi</i> (Latreille)	L+	C+
ARANEIDAE		
<i>Gibbaranea gibbosa</i> (Walckenaer)	-	C
<i>Araneus diadematus</i> Clerck	L+	C+
<i>Araneus quadratus</i> Clerck	L+	C+
<i>Atca sturmi</i> (Hahn)	L+	C
<i>Larinoides cornutus</i> (Clerck)	L+	C+
<i>Larinoides sclopetarius</i> (Clerck)	L+	C+
<i>Larinoides patagiatus</i> (Clerck)	-	C*
<i>Nuctenea umbratica</i> (Clerck)	L+	C+
<i>Agelenata redii</i> (Scopoli)	L+	C+
<i>Araniella cucurbitina</i> (Clerck)	L+	C+
<i>Araniella opistographa</i> (Kulczynski)	L+	C+
<i>Hyposinga pygmaea</i> (Sundevall)	L+	C+
<i>Cercidia prominens</i> (Westring)	-	C
<i>Zygiella x-notata</i> (Clerck)	L+	C+
<i>Zygiella atrica</i> (C.L. Koch)	L+	C+
<i>Mangora acalypha</i> (Walckenaer)	L*d	-
<i>Cyclosa conica</i> (Pallas)	L+	C+
LINYPHIIDAE		
<i>Ceratinella brevipes</i> (Westring)	L+	C+
<i>Ceratinella brevis</i> (Wider)	L+	C+
<i>Ceratinella scabrosa</i> (O.P.- Cambridge)	L	C+
<i>Walckenaeria dysderoides</i> (Wider)	-	C+
<i>Walckenaeria nudipalpis</i> (Westring)	L+	C+
<i>Walckenaeria vigilax</i> (Blackwall)	L+	C+
<i>Walckenaeria antica</i> (Wider)	L+	C+
<i>Walckenaeria cucullata</i> (C.L. Koch)	-	C+

<i>Walckenaeria nodosa</i> O.P.- Cambridge	L*	C+
<i>Walckenaeria incisa</i> (O.P.- Cambridge)	L+	-
<i>Walckenaeria cuspidata</i> (Blackwall)	L+	C+
<i>Walckenaeria furcillata</i> (Menge)	-	C+
<i>Walckenaeria atrotibialis</i> (O.P.- Cambridge)	L+	C
<i>Walckenaeria capito</i> (Westring)	L+	-
<i>Walckenaeria unicornis</i> O.P.- Cambridge	L+	C+
<i>Walckenaeria monoceros</i> (Wider)	L+	C+
<i>Walckenaeria kochi</i> (O.P.- Cambridge)	L+	C+
<i>Walckenaeria acuminata</i> Blackwall	L+	C+
<i>Dicymbium nigrum</i> (Blackwall)	L+	C+
<i>Dicymbium nigrum</i> f. <i>brevisetosum</i> Locket	L+	C+
<i>Dicymbium tibiale</i> (Blackwall)	L+	C+
<i>Entelecara flavipes</i> (Blackwall)	L	C+
<i>Entelecara erythropus</i> (Westring)	L+	C+
<i>Moebelia penicillata</i> (Westring)	L+	C+
<i>Hylaphantes graminicola</i> (Sundevall)	L+	C+
<i>Gnathonarium dentatum</i> (Wider)	L+	C+
<i>Tmeticus affinis</i> (Blackwall)	-	C+
<i>Gongylidium rufipes</i> (Sundevall)	L+	C+
<i>Dismodicus bifrons</i> (Blackwall)	L+	C+
<i>Hypomma bituberculatum</i> (Wider)	L+	C+
<i>Hypomma cornutum</i> (Blackwall)	L+	C+
<i>Metopobactrus prominulus</i> (O.P.- Cambridge)	L+	C+
<i>Baryphyma pratense</i> (Blackwall)	L	C+
<i>Baryphyma trifrons</i> (O.P.- Cambridge)	L+	C+
<i>Gonatium rubens</i> (Blackwall)	L+	C+
<i>Gonatium rubellum</i> (Blackwall)	L	C
<i>Maso sundevalli</i> (Westring)	L+	C+
<i>Peponocranium ludicum</i> (O.P.- Cambridge)	L+	C+
<i>Pocadicnemis pumila</i> (Blackwall)	L+	C+
<i>Pocadicnemis juncea</i> Locket & Millidge	L+	C+
<i>Hypselistes jacksoni</i> (O.P.- Cambridge)	-	C+
<i>Oedothorax gibbosus</i> (Blackwall)	L+	C+
<i>Oedothorax gibbosus</i> f. <i>tuberosus</i> (Blackwall)	L+	C+
<i>Oedothorax fuscus</i> (Blackwall)	L+	C+
<i>Oedothorax agrestis</i> (Blackwall)	L+	C+

<i>Oedothorax retusus</i> (Westring)	L+	C+
<i>Oedothorax apicatus</i> (Blackwall)	L+	C+
<i>Trichopterna thorelli</i> (Westring)	L	C
<i>Trichopterna mengei</i> (Simon)	-	C+
<i>Pelecopsis parallela</i> (Wider)	L+	C+
<i>Pelecopsis nemoralis</i> (Blackwall)	L+	C+
<i>Silometopus elegans</i> (O.P.- Cambridge)	L*	C+
<i>Silometopus ambiguus</i> (O.P.- Cambridge)	L+	C+
<i>Silometopus reussi</i> (Thorell)	-	C+
<i>Mecopisthes peusi</i> Wunderlich	L+	C+
<i>Cnephalocotes obscurus</i> (Blackwall)	L+	C+
<i>Ceratinopsis romana</i> (O.P.- Cambridge)	-	C+
<i>Evansia merens</i> O.P.- Cambridge	-	C+
<i>Tiso vagans</i> (Blackwall)	L+	C+
<i>Troxochrus scabriculus</i> (Westring)	L+	C+
<i>Troxochrus scabriculus</i> f. <i>cirrifrons</i> (O.P.- Cambridge)	L+	C+
<i>Minyriolus pusillus</i> (Wider)	L+	C+
<i>Tapinocyba praecox</i> (O.P.- Cambridge)	L+	C+
<i>Tapinocyba pallens</i> (O.P.- Cambridge)	L+	C+
<i>Tapinocyba insecta</i> (L. Koch)	L	-
<i>Microctenonyx subitaneus</i> (O.P.- Cambridge)	L	C+
<i>Satiliatas britteni</i> (Jackson)	L	C+
<i>Thyreosthenius parasiticus</i> (Westring)	L+	C+
<i>Monocephalus fuscipes</i> (Blackwall)	L+	C+
<i>Monocephalus castanicipes</i> (Simon)	L+	C+
<i>Lophomma punctatum</i> (Blackwall)	L+	C+
<i>Carorita limnaea</i> Crosby & Bishop	-	C+
<i>Gongylidiellum vivum</i> (O.P.- Cambridge)	L+	C+
<i>Gongylidiellum latebricola</i> (O.P.- Cambridge)	-	C
<i>Micrargus herbigradus</i> (Blackwall)	L+	C+
<i>Micrargus subaequalis</i> (Westring)	L+	C+
<i>Notioscopus sarcinatus</i> (O.P.- Cambridge)	-	C+
<i>Glypheais cottonae</i> (La Touche)	-	C+
<i>Erigonella hiemalis</i> (Blackwall)	L+	C+
<i>Erigonella ignobilis</i> (O.P.- Cambridge)	-	C+
<i>Savignya frontata</i> (Blackwall)	L+	C+
<i>Diplocephalus cristatus</i> (Blackwall)	L+	C+

<i>Diplocephalus permixtus</i> (O.P.- Cambridge)	L+	C+
<i>Diplocephalus latifrons</i> (O.P.- Cambridge)	L+	C+
<i>Diplocephalus picinus</i> (Blackwall)	L+	C+
<i>Araeoncus humilis</i> (Blackwall)	L+	C+
<i>Araeoncus crassiceps</i> (Westring)	L	C+
<i>Panamomops sulcifrons</i> (Wider)	-	C*
<i>Lessertia dentichelis</i> (Simon)	L+	C+
<i>Scotinotylus evansi</i> (O.P.- Cambridge)	L+	C+
<i>Typhocrestus digitatus</i> (O.P.- Cambridge)	L+	C+
<i>Milleriana inerrans</i> (O.P.- Cambridge)	L+	C+
<i>Erigone vagans</i> Audouin	L+	C+
<i>Erigone dentipalpis</i> (Wider)	L+	C+
<i>Erigone promiscua</i> (O.P.- Cambridge)	L+	C+
<i>Erigone atra</i> (Blackwall)	L+	C+
<i>Erigone arctica</i> (White)	L+	C+
<i>Erigone longipalpis</i> (Sundevall)	L+	C+
<i>Leptorhopitrum robustum</i> (Westring)	L+	C+
<i>Drepanotylus uncatus</i> (O.P.- Cambridge)	L+	C+
<i>Leptothrix hardyi</i> (Blackwall)	L+	C+
<i>Hilaira excisa</i> (O.P.- Cambridge)	L+	C+
<i>Halorates reprobus</i> (O.P.- Cambridge)	L+	C+
<i>Halorates distinctus</i> (Simon)	L+	C
<i>Carorita limnaea</i> Crosby & Bishop	-	C+
<i>Mioxena blanda</i> (Simon)	L+	C
<i>Astenargus paganus</i> (Simon)	-	C+
<i>Ostearius melanopygus</i> (O.P.- Cambridge)	L+	C+
<i>Aphileta misera</i> (O.P.- Cambridge)	L	C+
<i>Porrhomma pygmaeum</i> (Blackwall)	L+	C+
<i>Porrhomma convexum</i> (Westring)	L+	C+
<i>Porrhomma pallidum</i> Jackson	L+	C+
<i>Porrhomma campbelli</i> F.O.P.- Cambridge	L*	-
<i>Porrhomma errans</i> (Blackwall)	L*	-
<i>Porrhomma oblitum</i> (O.P.- Cambridge)	-	C+
<i>Porrhomma montanum</i> Jackson	L+	C
<i>Porrhomma microphthalmum</i> (O.P.- Cambridge)	-	C*
<i>Agyneta subtilis</i> (O.P.- Cambridge)	L+	C+
<i>Agyneta decora</i> (O.P.- Cambridge)	L+	C+

<i>Agyneta cauta</i> (O.P.- Cambridge)	L+	C+
<i>Agyneta olivacea</i> (Emerton)	-	C+
<i>Agyneta conigera</i> (O.P.- Cambridge)	L+	C+
<i>Agyneta ramosa</i> Jackson	L+	C
<i>Meioneta innotabilis</i> (O.P.- Cambridge)	L+	C+
<i>Meioneta rurestris</i> (C.L. Koch)	L+	C+
<i>Meioneta gulosa</i> (L. Koch)	L	C+
<i>Meioneta beata</i> (O.P.- Cambridge)	-	C*
<i>Meioneta mollis</i> (O.P.- Cambridge)	-	C+
<i>Meioneta saxatilis</i> (Blackwall)	L+	C+
<i>Microneta viaria</i> (Blackwall)	L+	C+
<i>Centromerus sylvaticus</i> (Blackwall)	L+	C+
<i>Centromerus prudens</i> (O.P.- Cambridge)	L+	C+
<i>Centromerus levitansis</i> (Simon)	-	C*
<i>Centromerus dilutus</i> (O.P.- Cambridge)	L+	L+
<i>Tallusia experta</i> (O.P.- Cambridge)	L+	C+
<i>Centromerita bicolor</i> (Blackwall)	L+	C+
<i>Centromerita concinna</i> (Thorell)	L+	C+
<i>Sintula cornigera</i> (Blackwall)	L*	C*
<i>Saaristoa abnormis</i> (Blackwall)	L+	C+
<i>Saaristoa firma</i> (O.P.- Cambridge)	L+	C+
<i>Macrargus rufus</i> (Wider)	L+	C+
<i>Bathyphantes approximatus</i> (O.P.- Cambridge)	L+	C+
<i>Bathyphantes gracilis</i> (Blackwall)	L+	C+
<i>Bathyphantes parvulus</i> (Westring)	L+	C+
<i>Bathyphantes nigrinus</i> (Westring)	L+	C+
<i>Bathyphantes setiger</i> F.O.P.- Cambridge	-	C+
<i>Kaestneria dorsalis</i> (Wider)	L+	C+
<i>Kaestneria pullata</i> (O.P.- Cambridge)	L+	C+
<i>Diplostyla concolor</i> (Wider)	L+	C+
<i>Poeciloneta globosa</i> (Wider)	L+	C+
<i>Drapetisca socialis</i> (Sundevall)	L+	C+
<i>Tapinopa longidens</i> (Wider)	L+	C+
<i>Floronia bucculenta</i> (Clerck)	L	C+
<i>Taranuncus setosus</i> (O.P.- Cambridge)	L	C
<i>Labulla thoracica</i> (Wider)	L+	C+
<i>Stemonyphantes lineatus</i> (Linnæus)	L+	C+

<i>Bolyphantes luteolus</i> (Blackwall)	L+	C+
<i>Bolyphantes alticeps</i> (Sundevall)	L+	C+
<i>Leptophantes nebulosus</i> (Sundevall)	L+	C+
<i>Leptophantes leprosus</i> (Ohlert)	L+	C+
<i>Leptophantes minutus</i> (Blackwall)	L+	C+
<i>Leptophantes alacris</i> (Blackwall)	L+	C+
<i>Leptophantes obscurus</i> (Blackwall)	L+	C+
<i>Leptophantes tenuis</i> (Blackwall)	L+	C+
<i>Leptophantes zimmermanni</i> Bertkau	L+	C+
<i>Leptophantes cristatus</i> (Menge)	L+	C+
<i>Leptophantes mengei</i> Kulczynski	L+	C+
<i>Leptophantes flavipes</i> (Blackwall)	L+	C+
<i>Leptophantes tenebricola</i> (Wider)	L+	C+
<i>Leptophantes ericaeus</i> (Blackwall)	L+	C+
<i>Leptophantes pallidus</i> (O.P.- Cambridge)	L+	C+
<i>Leptophantes insignis</i> O.P.- Cambridge	L+	C
<i>Helophora insignis</i> (Blackwall)	L+	C+
<i>Linyphia triangularis</i> (Clerck)	L+	C+
<i>Linyphia hortensis</i> Sundevall	L+	C+
<i>Linyphia montana</i> (Clerck)	L+	C+
<i>Linyphia clathrata</i> (Sundevall)	L+	C+
<i>Linyphia peltata</i> (Wider)	L+	C+
<i>Microlinyphia pusilla</i> (Sundevall)	L+	C+
<i>Microlinyphia impigra</i> (O.P.- Cambridge)	-	C+
<i>Allomengea scopigera</i> (Grube)	L+	C+
<i>Allomengea vidua</i> (L. Koch)	L+	C+

Acknowledgements:

I would like to thank Steve Judd for his advice on preparing this List, James Wright, Stan Dobson and Alan Scott for their help with records, and also Tony Parker for his kind assistance with computer documentation.

References:

- Blackwall, J.** (1861). A History of the Spiders of Great Britain and Ireland, Ray Society, London.
- Bristowe, W.S.** (1939). The Comity of Spiders, Ray Society, Vol. 1, London.
- Falconer, W.** (1930). Checklist of Lancashire and Cheshire Spiders. The Lancashire & Cheshire Fauna Committee, Pt. 1. pp. 81 - 89.
- Lockett, G.A. & Millidge, A.F.** (1951 & 1953). British Spiders, Ray Society, Vols. 1 & 2, London.
- Mackie, D.W.** (1960-77). Reports on Spiders. Lancashire and Cheshire Fauna Society Reports.
- Mackie, D.W.** (1975). The H.W. Freiston Collection of Spiders and Harvestmen in the Manchester Museum. B.A.S. Newsletter, 12 : 5-6.
- Roberts, M.J.** (1985). The Spiders of Great Britain and Ireland, Vols. 1 & 3, Harley Books, Colchester.
- Roberts, M.J.** (1987). The Spiders of Great Britain and Ireland, Vol. 2, Harley Books, Colchester.