

A Provisional Checklist to the Curculionoidea of Cheshire (VC58), South Lancashire (VC59)
and West Lancashire (VC60).
v.1.0, 25/2/2020

Clive Washington

This checklist covers all species in the Curculionoidea (currently 634 U.K. species as of 2020) in Cheshire, South Lancashire, and West Lancashire. It is derived in most part from data supplied to the National Curculionoidea Recording Scheme organised by Adrian Fowles. One of the by-products of this database is that it enables the relatively straightforward (!) production of county checklists, and Adrian has been so kind as to supply the data from the scheme for this purpose. The data have been provided by a wide range of individual recorders, county recorders, local record centres, and has been extracted from the literature and various collections. To this I have added some very recent records from my own efforts, as well as those from colleagues active in the area, which have not yet filtered through to the national scheme.

It would be quite clearly erroneous to suggest that any checklist was ‘complete’ in any way. There are always sources of records which have not been obtained – either published obscurely or never revealed at all. In the present case there are still some historical data sources which need further examination. However, the decision has been made to release the checklist in its present state, for a number of reasons. Firstly, because some of the old data may be very time-consuming to extract. Additionally, the appearance of a checklist often has the effect of stimulating the release of other records that were hidden in the woodwork (perhaps literally in the case of weevils!). And finally, it stimulates ongoing recording efforts, allowing recorders to focus on gaps in the knowledge base and gain some idea of the scarcity of the species they find. The expectation is that the list will be revised on a fairly regular basis as new data emerges.

Individual entries in the checklist are intended to give some indication as to the currency of the information, by providing the number of records in the recording scheme, and the year of the most recent record. For the more commonly recorded species – generally those with more than 8-10 records on the database – I have simply indicated them as ‘Frequent’. In some cases experience shows that they are often found in quantity, and in these cases the designation has been changed to ‘Abundant’. In both cases this should be taken to mean “in the appropriate habitat”. So for example, whereas *Euophryum confine* is almost ubiquitous in dead wood, you would struggle to find it on sand dunes.

Individual records, dates, recorders, and so on, have not been included for several reasons. They would of course bloat the document. But primarily they have been omitted since some data was originally supplied to the National Recording Scheme under various restrictions; sometimes out of privacy concerns, or due to a desire to avoid duplication, or as a result of specific licensing or contractual issues. If anyone should require further information - if they are seeking a particular species, for example – then it is often possible for myself, or the National Recording Scheme, to provide more detail in many cases.

It is interesting to compare the relative records across the three vice-counties. Cheshire has records for 275 species, South Lancashire 231 species, and West Lancashire only 129 species. This is partly due to differences in the diversity of habitats but also reflects the recording efforts of the past. Cheshire has a diverse mix of grassland, meadow, wetland, moss, and woodland, and includes the Wirral and its coastal sites. South Lancashire includes a large stretch of the Sefton Coast which has historically provided records for many rare Coleoptera. West Lancashire also includes coastal regions, as well as upland and forested areas, but the recording scheme only has four records for *Sitona lineatus*, probably the most abundant of all weevils! This must surely give an indication that recording effort in VC60 has been significantly less than in the other vice-counties, and that much work remains to be done in this region.

| | Preferred Species Name | VC58 Cheshire | VC59 South Lancashire | VC60 West Lancashire |
|------------------------------|--|------------------------------------|-----------------------------|-----------------------------|
| Nemonchyidae (1 sp.) | | | | |
| | <i>Cimberis attelaboides</i> (Fabricius, 1787) | Single record 1908 | | |
| Anthribidae (9 sp.) | | | | |
| | <i>Anthribus fasciatus</i> Forster, 1770 | 2 records, 1908 & 1991 | | |
| | <i>Anthribus nebulosus</i> Forster, 1770 | Single record 1908 | | |
| | <i>Platyrhinus resinosus</i> (Scopoli, 1763) | Single record 2019 | Single record 1992 | |
| | <i>Platystomos albinus</i> (Linnaeus, 1758) | | Single record 1993 | |
| | <i>Choragus sheppardi</i> Kirby, 1819 | 4 records, most recent 1992 | Single record 2015 | Single record 1993 |
| Rhynchitidae (19 sp.) | | | | |
| | <i>Involvulus cupreus</i> (Linnaeus, 1758) | 3 records, most recent 2019 | | |
| | <i>Involvulus icosandriae</i> (Scopoli, 1763) | | Single record 2009 | |
| | <i>Lasiorrhynchites cavifrons</i> (Gyllenhal, 1833) | Single record 2017 | | |
| | <i>Neocoenorrhinus germanicus</i> (Herbst, 1797) | Single record pre-1900 | Single record 2011 | Single record 1999 |
| | <i>Neocoenorrhinus interpunctatus</i> (Stephens, 1831) | Single record 1961 | | |
| | <i>Neocoenorrhinus minutus</i> (Herbst, 1797) | 3 records, most recent 1970 | | |
| | <i>Tatianaerhynchites aequatus</i> (Linnaeus, 1767) | Frequent across Vice-County | | Single record 2006 |
| | <i>Temnocerus longiceps</i> (C. G. Thomson, 1888) | 2 records, 1990 & 2004 | 3 records, most recent 2011 | Single record 2018 |
| | <i>Temnocerus namus</i> (Paykull, 1792) | Frequent across Vice-County | Frequent across Vice-County | |
| | <i>Caenorhinus mannerheimii</i> (Hummel, 1823) | 8 records, most recent 2000 | 3 records, most recent 2000 | |
| | <i>Deporaus betulae</i> (Linnaeus, 1758) | Abundant across Vice-County | Frequent across Vice-County | Single record 1999 |
| Attelabidae (2 sp.) | | | | |
| | <i>Attelabus nitens</i> (Scopoli, 1763) | 6 records, most recent 1995 | Single record 2015 | |
| | <i>Apoderus coryli</i> (Linnaeus, 1758) | 3 records, most recent 2018 | 9 records, most recent 2017 | |
| Apionidae (90 sp.) | | | | |
| | <i>Apion cruentatum</i> Walton, 1844 | 2 records pre-1900 and one in 2020 | | |
| | <i>Apion frumentarium</i> (Linnaeus, 1758) | Abundant across Vice-County | Frequent across Vice-County | 2 records, most recent 2017 |
| | <i>Apion haematodes</i> Kirby, 1808 | 5 records, most recent 2015 | 3 records, most recent 2015 | |
| | <i>Apion rubens</i> Walton, 1837 | Frequent across Vice-County | Single record 1997 | |
| | <i>Apion rubiginosum</i> Grill, 1893 | Single record 2008 | | |
| | <i>Perapion affine</i> (Kirby, 1808) | 2 records, 1917 & 1966 | 2 records, 1960 & 1991 | Single record 1950 |
| | <i>Perapion curtirostre</i> (Germar, 1817) | Frequent across Vice-County | 3 records, most recent 2016 | 4 records, 2017 and 2018 |
| | <i>Perapion hydrolapathi</i> (Marshall, 1802) | Abundant across Vice-County | Frequent across Vice-County | Single record 2018 |
| | <i>Perapion marchicum</i> (Herbst, 1797) | Single record 2004 | 4 records, most recent 2007 | Single record 2014 |
| | <i>Perapion violaceum</i> (Kirby, 1808) | Frequent across Vice-County | 3 records, most recent 2007 | 2 records, 2018 |
| | <i>Pseudaplemonus limonii</i> (Kirby, 1808) | | | 6 records, 1991 and 2018 |
| | <i>Aspidapion radiolus</i> (Marshall, 1802) | 4 records, most recent 2013 | | |
| | <i>Aspidapion aeneum</i> (Fabricius, 1775) | Single record 1991 | | |

| Preferred Species Name | VC58 Cheshire | VC59 South Lancashire | VC60 West Lancashire |
|--|-----------------------------|------------------------------|-----------------------------|
| <i>Ceratapion carduorum</i> (Kirby, 1808) | 5 records, most recent 2016 | 2 records, most recent 2010 | |
| <i>Ceratapion gibbirostre</i> (Gyllenhal, 1813) | 9 records, most recent 2016 | 6 records, most recent 2018 | |
| <i>Ceratapion onopordi</i> (Kirby, 1808) | Frequent across Vice-County | 5 records, most recent 2017 | |
| <i>Diplapion confluens</i> (Kirby, 1808) | 2 records, most recent 2016 | | |
| <i>Omphalapion hookerorum</i> (Kirby, 1808) | Single record 2016 | Single record 2017 | |
| <i>Exapion difficile</i> (Herbst, 1797) | | | Single record 1950 |
| <i>Exapion fuscirostre</i> (Fabricius, 1775) | Single record 2001 | Single record 2016 | Single record 2008 |
| <i>Exapion ulicis</i> (Forster, 1771) | Abundant across Vice-County | Abundant across Vice-County | Single record 2016 |
| <i>Kalcapion pallipes</i> (Kirby, 1808) | Single record 1999 | | |
| <i>Taeniapion urticarium</i> (Herbst, 1784) | | Single record 1989 | |
| <i>Malvapion malvae</i> (Fabricius, 1775) | 3 records, most recent 2004 | | |
| <i>Pseudapion rufirostre</i> (Fabricius, 1775) | 2 records, most recent 2004 | Single record 1991 | |
| <i>Cyanapion spencii</i> (Kirby, 1808) | | 8 records, most recent 2010 | Single record 1975 |
| <i>Cyanapion afer</i> (Gyllenhal, 1833) | Single record pre-1900 | | |
| <i>Cyanapion gyllenhalii</i> (Kirby, 1808) | Single record 2010 | | Single record 1975 |
| <i>Eutrichapion ervi</i> (Kirby, 1808) | 9 records, most recent 2018 | Frequent across Vice-County | 3 records, most recent 2016 |
| <i>Eutrichapion viciae</i> (Paykull, 1800) | 4 records, most recent 2008 | 10 records, most recent 2017 | Single record 1975 |
| <i>Eutrichapion vorax</i> (Herbst, 1797) | 4 records, most recent 2015 | | |
| <i>Eutrichapion punctigerum</i> (Paykull, 1792) | Single record 1934 | | |
| <i>Holotrichapion ononis</i> (Kirby, 1808) | Single record 2016 | 5 records, most recent 2009 | |
| <i>Holotrichapion pisi</i> (Fabricius, 1801) | Single record 2016 | | Single record 2006 |
| <i>Holotrichapion aethiops</i> (Herbst, 1797) | 2 records 1909 & 2013 | 2 records, 2010 & 2015 | |
| <i>Oxystoma cerdo</i> (Gerstäcker, 1854) | 8 records, most recent 2016 | 4 records, most recent 2016 | 3 records, most recent 2016 |
| <i>Oxystoma cracca</i> (Linnaeus, 1767) | 2 records, most recent 2016 | Single record 1999 | |
| <i>Oxystoma pomonae</i> (Fabricius, 1798) | 7 records, most recent 2015 | Abundant coastally | |
| <i>Oxystoma subulatum</i> (Kirby, 1808) | 3 records, most recent 2013 | Frequent across Vice-County | |
| <i>Pirapion immune</i> (Kirby, 1808) | Single record 1980 | Single record 1945 | |
| <i>Catapion pubescens</i> (Kirby, 1811) | | Single record 2018 | |
| <i>Catapion seniculus</i> (Kirby, 1808) | Single record 1963 | 2 records, 1985 & 2017 | |
| <i>Ischnopterapion loti</i> (Kirby, 1808) | Abundant across Vice-County | Abundant across Vice-County | 2 records, 1993 & 2008 |
| <i>Ischnopterapion modestum</i> (Germar, 1817) | Single record 2001 | Single record 2017 | Single record 2016 |
| <i>Ischnopterapion virens</i> (Herbst, 1797) | Frequent across Vice-County | 5 records, most recent 2018 | Single record 2016 |
| <i>Protopirapion atratum</i> (Germar, 1817) | Single record 2009 | | Single record 2014 |
| <i>Stenopterapion tenue</i> (Kirby, 1808) | 2 records, 2016 & 2019 | | |
| <i>Stenopterapion scutellare</i> (Kirby, 1811) | 2 records, 1908 & 2017 | | |
| <i>Betulapion simile</i> (Kirby, 1811) | 3 records, most recent 2010 | Frequent across Vice-County | 3 records, most recent 2017 |
| <i>Protapion apricans</i> (Herbst, 1797) | Frequent across Vice-County | Frequent across Vice-County | Frequent across Vice-County |
| <i>Protapion assimile</i> (Kirby, 1808) | 5 records, most recent 2016 | 5 records, most recent 2017 | 2 records, most recent 2016 |
| <i>Protapion diffforme</i> (Germar, 1818) | Single record 2013 | | |
| <i>Protapion dissimile</i> (Germar, 1817) | Single record 2016 | | |
| <i>Protapion fulvipes</i> (Geoffroy in Fourcroy, 1785) | Abundant across Vice-County | Abundant across Vice-County | 2 records, most recent 2014 |
| <i>Protapion nigrirtarse</i> (Kirby, 1808) | 2 records pre-1900 | 2 records, 1985 & 2007 | Single record 1975 |
| <i>Protapion ononidis</i> (Gyllenhal, 1827) | 4 records, most recent 2015 | 9 records, most recent 2005 | |
| <i>Protapion trifolii</i> (Linnaeus, 1768) | 6 records, most recent 2016 | 9 records, most recent 2018 | 3 records, most recent 2017 |

| | Preferred Species Name | VC58 Cheshire | VC59 South Lancashire | VC60 West Lancashire |
|--|---|-----------------------------|------------------------------|-----------------------------|
| | | | | |
| | Nanophyidae (2 sp.) | | | |
| | <i>Nanophyes marmoratus</i> (Goeze, 1777) | 6 records, most recent 2015 | 6 records, most recent 2018 | |
| | | | | |
| | Dryophthoridae (4 sp) | | | |
| | <i>Sitophilus oryzae</i> (Linnaeus, 1763) | 2 records, 1981 & 2011 | Single record 2005 | |
| | <i>Sitophilus zeamais</i> Motschulsky, 1855 | Single record 2003 | | |
| | | | | |
| | Eirrhinidae (13 sp) | | | |
| | <i>Grypus equiseti</i> (Fabricius, 1775) | Frequent across Vice-County | Frequent across Vice-County | |
| | <i>Notaris acridulus</i> (Linnaeus, 1758) | Frequent across Vice-County | 2 records, 1990 & 2012 | Single record 2014 |
| | <i>Notaris scirpi</i> (Fabricius, 1792) | | 9 records, most recent 2015 | |
| | <i>Thryogenes festucae</i> (Herbst, 1795) | 2 records 1991 | | |
| | <i>Thryogenes nereis</i> (Paykull, 1800) | 6 records, most recent 2015 | 2 records, 2011 | |
| | <i>Tournotaris bimaculatus</i> (Fabricius, 1787) | 2 records, most recent 2005 | 5 records, most recent 2005 | Single record 2017 |
| | <i>Stenopelmus rufinasus</i> Gyllenhal in Schönherr, 1835 | 6 records, most recent 2016 | 3 records, most recent 2018 | |
| | <i>Tanysphyrus lemnae</i> (Paykull, 1792) | Frequent across Vice-County | Single record 2018 | |
| | | | | |
| | Raymondionymidae (1 sp.) – Not yet recorded in VC58-60 | | | |
| | | | | |
| | Curculionidae (493 sp.) | | | |
| | Curculioninae | | | |
| | <i>Archarius pyrrhoceras</i> (Marsham, 1802) | Abundant across Vice-County | 6 records, most recent 2012 | 2 records, 2011 |
| | <i>Archarius salicivorus</i> (Paykull, 1792) | Abundant across Vice-County | Frequent across Vice-County | 3 records, most recent 2017 |
| | <i>Curculio betulae</i> (Stephens, 1831) | | Single record 1978 | Single record 2017 |
| | <i>Curculio glandium</i> Marsham, 1802 | 6 records, most recent 2013 | 6 records, most recent 2013 | |
| | <i>Curculio nucum</i> Linnaeus, 1758 | 2 records, most recent 2009 | Single record 2016 | |
| | <i>Curculio rubidus</i> (Gyllenhal, 1835) | Single record 2017 | Single record 2013 | |
| | <i>Curculio venosus</i> (Gravenhorst, 1807) | 9 records, most recent 2012 | | |
| | <i>Curculio villosus</i> Fabricius, 1781 | 2 records, most recent 2009 | Single record 1987 | |
| | <i>Anthonomus pedicularius</i> (Linnaeus, 1758) | 7 records, most recent 2016 | 7 records, most recent 2017 | 2 records, 2011 |
| | <i>Anthonomus pomorum</i> (Linnaeus, 1758) | Single record 2009 | Single record 2013 | 3 records, most recent 2011 |
| | <i>Anthonomus rubi</i> (Herbst, 1795) | 7 records, most recent 2016 | 3 records, most recent 2016 | 2 records, 2016 |
| | <i>Anthonomus ulmi</i> (De Geer, 1775) | | Single record 2011 | |
| | <i>Cionus alauda</i> (Herbst, 1784) | | Single record pre-1900 | |
| | <i>Cionus hortulanus</i> (Geoffroy in Fourcroy, 1785) | | 3 records, most recent 2015 | |
| | <i>Cionus scrophulariae</i> (Linnaeus, 1758) | Abundant across Vice-County | Abundant across Vice-County | 4 records, most recent 2018 |
| | <i>Dorytomus dejeani</i> Faust, 1883 | 3 records, most recent 2013 | 6 records, most recent 2017 | |
| | <i>Dorytomus longimanus</i> (Forster, 1771) | 2 records, 1992 & 2018 | 4 records, most recent 1996 | |
| | <i>Dorytomus majalis</i> (Paykull, 1792) | Single record 1949 | | |
| | <i>Dorytomus melanophthalmus</i> (Paykull, 1792) | | Single record 1995 | |
| | <i>Dorytomus rufatus</i> (Bedel, 1888) | 6 records, most recent 2010 | Single record 2009 | Single record 2016 |
| | <i>Dorytomus taeniatus</i> (Fabricius, 1781) | 7 records, most recent 2016 | 10 records, most recent 2012 | 3 records, most recent 2017 |

| Preferred Species Name | VC58 Cheshire | VC59 South Lancashire | VC60 West Lancashire |
|---|-----------------------------|-----------------------------|-----------------------------|
| <i>Dorytomus tortrix</i> (Linnaeus, 1760) | | 5 records, most recent 2016 | |
| <i>Dorytomus tremulae</i> (Fabricius, 1787) | | 2 records, 1963 | |
| <i>Gymnetron beccabungae</i> s.lat. | Single record 1980 | | |
| <i>Gymnetron veronicae</i> (Germar, 1821) | Single record 2009 | | |
| <i>Mecinus circulatus</i> (Marshall, 1802) | Single record 1991 | | |
| <i>Mecinus collaris</i> Germar, 1821 | | Single record 1927 | |
| <i>Mecinus labilis</i> (Herbst, 1795) | | | Single record 1993 |
| <i>Mecinus pascuorum</i> (Gyllenhal, 1813) | Frequent across Vice-County | Frequent across Vice-County | 2 records, most recent 2017 |
| <i>Mecinus pyrastrer</i> (Herbst, 1795) | 9 records, most recent 2018 | 7 records, most recent 2016 | Single record 2014 |
| <i>Rhinusa antirrhini</i> (Paykull, 1800) | | Single record 1989 | |
| <i>Rhinusa collina</i> (Gyllenhal, 1813) | | Single record 1902 | |
| <i>Rhinusa linariae</i> (Panzer, 1795) | | 3 records, 1902-3 | |
| <i>Isochnus foliorum</i> (O. F. Müller, 1764) | 2 records, 1992 and 2020 | | |
| <i>Isochnus sequensi</i> (Steirlin, 1894) | | Single record 2018 | |
| <i>Orchestes alni</i> (Linnaeus, 1758) | Single record 2001 | Single record 2000 | |
| <i>Orchestes calceatus</i> (Germar, 1821) | Single record pre-1900 | | |
| <i>Orchestes hortorum</i> (Fabricius, 1792) | Single record 2012 | 4 records, most recent 2018 | |
| <i>Orchestes pilosus</i> (Fabricius, 1781) | 3 records, most recent 2009 | | |
| <i>Orchestes quercus</i> (Linnaeus, 1758) | Frequent across Vice-County | 4 records, most recent 2012 | |
| <i>Orchestes rusci</i> (Herbst, 1795) | Frequent across Vice-County | 4 records, most recent 2016 | 2 records, most recent 2011 |
| <i>Orchestes testaceus</i> (O. F. Müller, 1776) | 2 records, 1908 & 2009 | | |
| <i>Orchestes fagi</i> (Linnaeus, 1758) | Abundant across Vice-County | Abundant across Vice-County | 5 records, most recent 2018 |
| <i>Rhamphus oxyacanthae</i> (Marshall, 1802) | Frequent across Vice-County | | |
| <i>Rhamphus pulicarius</i> (Herbst, 1795) | Frequent across Vice-County | 7 records, most recent 2011 | 4 records, most recent 2011 |
| <i>Tachyerges salicis</i> (Linnaeus, 1758) | Single record 1991 | 5 records, most recent 2000 | |
| <i>Tachyerges stigma</i> (Germar, 1821) | 4 records, most recent 2004 | 6 records, most recent 2015 | Single record 1999 |
| <i>Orthochaetes setiger</i> (Beck, 1817) | 3 records, most recent 2019 | 5 records, most recent 2000 | 2 records, most recent 2019 |
| <i>Sibinia primita</i> (Herbst, 1795) | Single record 2015 | | |
| <i>Tychius junceus</i> (Reich, 1797) | | Single record 1965 | |
| <i>Tychius meliloti</i> Stephens, 1831 | Single record 2005 | | |
| <i>Tychius picirostris</i> (Fabricius, 1787) | Frequent across Vice-County | Frequent across Vice-County | Single record 2016 |
| Bagoinae | | | |
| <i>Bagous limosus</i> (Gyllenhal, 1827) | | 2 records, 1900 & 1995 | |
| <i>Bagous glabrirostris</i> (Herbst, 1795) | Single record 1997 | | |
| <i>Bagous lutosus</i> (Gyllenhal, 1813) | | 7 records, most recent 2019 | |
| <i>Bagous lutulentus</i> (Gyllenhal, 1813) | | 7 records, most recent 2018 | |
| Baridinae | | | |
| <i>Hydronomus alismatis</i> (Marshall, 1802) | 4 records, most recent 1991 | Single record pre-1900 | Single record 2006 |
| <i>Limnobaris dolorosa</i> (Goeze, 1777) | Single record 2004 | 3 records, most recent 2018 | 2 records, 2017 |
| <i>Limnobaris t-album</i> (Linnaeus, 1758) | 2 records, most recent 1982 | Single record 1990 | Single record 2013 |

| | Preferred Species Name | VC58 Cheshire | VC59 South Lancashire | VC60 West Lancashire |
|--|--|-----------------------------|-----------------------------|-----------------------------|
| | Ceutorhynchinae | | | |
| | <i>Calosirus terminatus</i> (Herbst, 1795) | Single record 2019 | Single record 1908 | |
| | <i>Ceutorhynchus alliariae</i> Brisout de Barneville, H., 1860 | | | Single record 1988 |
| | <i>Ceutorhynchus assimilis</i> (Paykull, 1792) | 4 records, most recent 2001 | 6 records, most recent 2010 | |
| | <i>Ceutorhynchus atomus</i> Boheman, 1845 | Single record 1950 | | |
| | <i>Ceutorhynchus cochleariae</i> (Gyllenhal, 1813) | | 2 records, 2010 & 2011 | 2 records, 2017 |
| | <i>Ceutorhynchus constrictus</i> (Marsham, 1802) | Single record 1992 | | |
| | <i>Ceutorhynchus contractus</i> (Marsham, 1802) | 6 records, most recent 2015 | Single record 2010 | |
| | <i>Ceutorhynchus erysimi</i> (Fabricius, 1787) | 7 records, most recent 2009 | 5 records, most recent 2015 | |
| | <i>Ceutorhynchus hirtulus</i> Germar, 1824 | | Single record 1977 | |
| | <i>Ceutorhynchus obstructus</i> (Marsham, 1802) | Single record 2008 | 4 records, most recent 2018 | 2 records, most recent 2015 |
| | <i>Ceutorhynchus pallidactylus</i> (Marsham, 1802) | 3 records, most recent 2000 | 5 records, most recent 2017 | |
| | <i>Ceutorhynchus pectoralis</i> Weise, 1895 | Single record 1991 | | |
| | <i>Ceutorhynchus picitarsis</i> Gyllenhal, 1837 | Single record 2014 | | |
| | <i>Ceutorhynchus pulvinatus</i> Gyllenhal, 1837 | Uncertain record 2011 | | |
| | <i>Ceutorhynchus pumilio</i> (Gyllenhal, 1827) | | | Single record pre-1900 |
| | <i>Ceutorhynchus querceti</i> (Gyllenhal, 1813) | Uncertain record 1979 | | |
| | <i>Ceutorhynchus rapae</i> Gyllenhal, 1837 | Single record 2012 | | |
| | <i>Ceutorhynchus typhae</i> (Herbst, 1795) | 7 records, most recent 2018 | 2 records, 2009 & 2016 | 2 records, 1993 & 2016 |
| | <i>Coeliodes rana</i> (Fabricius, 1787) | 2 records, 2009 & 2010 | | 3 records 2011 |
| | <i>Coeliodinus rubicundus</i> (Herbst, 1795) | 7 records, most recent 2014 | 3 records, most recent 2016 | Single record 1994 |
| | <i>Datonychus angulosus</i> (Boheman, 1845) | | Single record pre-1900 | |
| | <i>Datonychus arquata</i> (Herbst, 1795) | | 5 records pre-1900 | |
| | <i>Datonychus melanostictus</i> (Marsham, 1802) | 3 records, most recent 1994 | 2 records, 2008 & 2016 | |
| | <i>Glocianus distinctus</i> (Brisout de Barneville, C., 1870) | | Single record 2017 | |
| | <i>Glocianus punctiger</i> (Sahlberg, C.R., 1835) | Single record 1939 | Single record 2018 | |
| | <i>Hadroplontus litura</i> (Fabricius, 1775) | Frequent across Vice-County | 8 records, most recent 2017 | |
| | <i>Micrelus ericae</i> (Gyllenhal, 1813) | 7 records, most recent 2014 | Frequent across Vice-County | Single record 2017 |
| | <i>Microplontus campestris</i> (Gyllenhal, 1837) | | Single record 1975 | |
| | <i>Microplontus melanostigma</i> (Marsham, 1802) | | Single record 1991 | |
| | <i>Nedyus quadrimaculatus</i> (Linnaeus, 1758) | Abundant across Vice-County | Abundant across Vice-County | 9 records, most recent 2017 |
| | <i>Parathelcus pollinarius</i> (Forster, 1771) | Abundant across Vice-County | Abundant across Vice-County | 2 records, 1975 & 2014 |
| | <i>Poophagus sisymbrii</i> (Fabricius, 1777) | 4 records, most recent 1991 | | Single record 2010 |
| | <i>Sirocalodes depressicollis</i> (Gyllenhal, 1813) | | Single record 1917 | |
| | <i>Sirocalodes mixtus</i> (Mulsant & Rey, 1859) | Single record 1991 | | |
| | <i>Thamiocolus viduatus</i> (Gyllenhal, 1813) | | 2 records, 1998 | |
| | <i>Trichosirocalus dawsoni</i> (Brisout de Barneville, C., 1869) | | | Single record 1989 |
| | <i>Trichosirocalus troglodytes</i> (Fabricius, 1787) | Frequent across Vice-County | Frequent across Vice-County | 5 records, most recent 2017 |
| | <i>Zacladus geranii</i> (Paykull, 1800) | 9 records, most recent 2015 | Frequent across Vice-County | Single record 1993 |
| | <i>Amalus scortillum</i> (Herbst, 1795) | 2 records, 2010 & 2016 | 2 records, 2017 | |
| | <i>Pelenomus commari</i> (Panzer, 1795) | 3 records, most recent 1991 | | |
| | <i>Pelenomus quadrituberculatus</i> (Fabricius, 1787) | 6 records, most recent 2014 | 4 records, most recent 2014 | |
| | <i>Pelenomus waltoni</i> (Boheman, 1843) | | | Single record 2011 |

| Preferred Species Name | VC58 Cheshire | VC59 South Lancashire | VC60 West Lancashire |
|--|-----------------------------|-----------------------------|-----------------------------|
| <i>Pelenomus zumpti</i> (Wagner, 1939) | Single record 1996 | | |
| <i>Phytobius leucogaster</i> (Marshall, 1802) | 3 records, most recent 1992 | | |
| <i>Rhinoncus bruchoides</i> (Herbst, 1784) | 2 records, 1963 and 1990 | 2 records, 2012 & 2017 | |
| <i>Rhinoncus castor</i> (Fabricius, 1792) | Frequent across Vice-County | Frequent across Vice-County | Single record 2017 |
| <i>Rhinoncus inconspicuous</i> (Herbst, 1795) | Single record 2016 | Single record 2017 | Single record 2010 |
| <i>Rhinoncus pericarpus</i> (Linnaeus, 1758) | Abundant across Vice-County | Abundant across Vice-County | 4 records, most recent 2017 |
| <i>Rhinoncus perpendicularis</i> (Reich, 1797) | Frequent across Vice-County | 5 records, most recent 2016 | |
| Cossoninae | | | |
| <i>Cossonus parallelepipeds</i> (Herbst, 1795) | Single record 1986 | | |
| <i>Rhopalomesites tardii</i> (Curtis, 1825) | 4 records pre-1900 | | |
| <i>Euophryum confine</i> (Broun, 1881) | Abundant across Vice-County | Abundant across Vice-County | 5 records, most recent 2017 |
| <i>Pentarthrum huttoni</i> Wollaston, 1854 | Single record 2015 | | |
| <i>Pentarthrum huttoni</i> Wollaston, 1854 | | 3 records, most recent 2012 | |
| <i>Phloeophagus lignarius</i> (Marshall, 1802) | 6 records, most recent 1990 | Single record pre-1900 | |
| Cryptorhynchinae | | | |
| <i>Cryptorhynchus lapathi</i> (Linnaeus, 1758) | Single record 1945 | 2 records, 1959 & 1987 | 2 records, pre-1900 & 2006 |
| <i>Acalles misellus</i> Boheman, 1844 | Single record 2018 | Single record 1994 | |
| <i>Acalles ptinoides</i> (Marshall, 1802) | 3 records, most recent 2018 | | Single record 1995 |
| Entiminae | | | |
| <i>Graptus triguttatus</i> (Fabricius, 1775) | 2 records, 1985 and 1993 | | |
| <i>Strophosoma capitatum</i> (De Geer, 1775) | Single record 1974 | Single record 1997 | |
| <i>Strophosoma faber</i> (Herbst, 1784) | Single record 1960 | | Single record 1950 |
| <i>Strophosoma melanogrammum</i> (Forster, 1771) | Abundant across Vice-County | Abundant across Vice-County | Abundant across Vice-County |
| <i>Strophosoma nebulosum</i> (Stephens, 1831) | Single record 2016 | | |
| <i>Strophosoma nebulosum</i> (Stephens, 1831) | | | 2 records, 1993 & 2005 |
| <i>Strophosoma sus</i> (Stephens, 1831) | 5 records, most recent 1991 | | Single record 1950 |
| <i>Philopeda plagiata</i> (Schaller, 1783) | Frequent in sandy areas | Frequent in sandy areas | 3 records, most recent 2010 |
| <i>Barynotus moerens</i> (Fabricius, 1792) | 2 records, 1985 and 2015 | Single record 2018 | |
| <i>Barynotus obscurus</i> (Fabricius, 1775) | Frequent across Vice-County | 6 records, most recent 2016 | |
| <i>Barynotus squamosus</i> Germar, 1824 | 2 records, 1979 & 2000 | Single record 1991 | |
| <i>Barynotus squamosus</i> Germar, 1824 | | | |
| <i>Otiorhynchus armadillo</i> (Rossi, 1792) | | Single record 2018 | Single record 2013 |
| <i>Otiorhynchus atroapterus</i> (De Geer, 1775) | | 5 records, most recent 2017 | |
| <i>Otiorhynchus crataegi</i> Germar, 1824 | | Single record 2013 | |
| <i>Otiorhynchus ligneus</i> (Olivier, 1807) | 2 records, 1979 & 1999 | | |
| <i>Otiorhynchus nodosus</i> (O. F. Müller, 1764) | | Single record pre-1900 | |
| <i>Otiorhynchus ovatus</i> (Linnaeus, 1758) | Frequent in sandy areas | Frequent in sandy areas | Single record 2010 |
| <i>Otiorhynchus porcatus</i> (Herbst, 1795) | Single record 2010 | | |
| <i>Otiorhynchus rugifrons</i> (Gyllenhal, 1813) | Single record 1979 | | Single record 2005 |
| <i>Otiorhynchus rugosostriatus</i> (Goeze, 1777) | 5 records, most recent 2010 | | |
| <i>Otiorhynchus singularis</i> (Linnaeus, 1767) | Frequent across Vice-County | Frequent across Vice-County | Frequent across Vice-County |

| Preferred Species Name | VC58 Cheshire | VC59 South Lancashire | VC60 West Lancashire |
|---|--------------------------------------|-----------------------------|----------------------------------|
| <i>Otiorynchus sulcatus</i> (Fabricius, 1775) | Frequent across Vice-County | Frequent across Vice-County | 2 records, 2011 |
| <i>Otiorynchus tenebricosus</i> (Herbst, 1784) | | 4 records, most recent 2006 | |
| <i>Phyllobius pyri</i> (Linnaeus, 1758) | Abundant across Vice-County | Abundant across Vice-County | Frequent across Vice-County |
| <i>Phyllobius argentatus</i> (Linnaeus, 1758) | Abundant across Vice-County | Abundant across Vice-County | Frequent across Vice-County |
| <i>Phyllobius glaucus</i> (Scopoli, 1763) | Abundant across Vice-County | 9 records, most recent 2018 | 2 records, 1994 & 2011 |
| <i>Phyllobius pomaceus</i> Gyllenhal, 1834 | Abundant across Vice-County | Abundant across Vice-County | 4 records, most recent 2018 |
| <i>Phyllobius oblongus</i> (Linnaeus, 1758) | Abundant across Vice-County | Abundant across Vice-County | 2 records, 2010 & 2011 |
| <i>Phyllobius roboretanus</i> Gredler, 1882 | Abundant across Vice-County | Abundant across Vice-County | 2 records, 2010 & 2018 |
| <i>Phyllobius viridicollis</i> (Fabricius, 1792) | 5 records, most recent 2015 | 8 records, most recent 2016 | 2 records, 1950 & 1999 |
| <i>Phyllobius maculicornis</i> Germar, 1824 | 7 records, most recent 2016 | | Single record 1950 |
| <i>Phyllobius virideaeris</i> (Laicharting, 1781) | 9 records, most recent 2015 | 5 records, most recent 2014 | 2 records, 2010 & 2017 |
| <i>Liophloeus tessulatus</i> (O. F. Müller, 1776) | 3 records, most recent 1991 | 6 records, most recent 2018 | Single record 2011 |
| <i>Polydrusus tereticollis</i> (De Geer, 1775) | 2 records, 2000 and 2006 | 2 records, 1995 & 1996 | 2 records, 2011 & 2017 |
| <i>Polydrusus formosus</i> (Mayer, 1779) | Frequent across Vice-County | Frequent across Vice-County | Single record 2017 |
| <i>Polydrusus mollis</i> (Ström, 1768) | 2 records, 1995 & 1996 | | Single record 2011 |
| <i>Polydrusus confluens</i> Stephens, 1831 | 2 uncertain records, pre-1900 & 1980 | | |
| <i>Polydrusus planifrons</i> Gyllenhal, 1834 | | | Single uncertain record pre-1900 |
| <i>Polydrusus pterygomalis</i> Boheman, 1840 | Frequent across Vice-County | Frequent across Vice-County | 7 records, most recent 2017 |
| <i>Polydrusus cervinus</i> (Linnaeus, 1758) | Frequent across Vice-County | Frequent across Vice-County | Single record 2011 |
| <i>Polydrusus pilosus</i> Gredler, 1866 | Single record 2003 | Single record 1936 | |
| <i>Polydrusus pulchellus</i> Stephens, 1831 | 3 records, most recent 1980 | 2 records, 1989 & 2005 | 3 records, most recent 2018 |
| <i>Exomias araneiformis</i> (Schrank, 1781) | 4 records, most recent 2016 | Frequent across Vice-County | 2 records, 2011 |
| <i>Exomias pellucidus</i> (Boheman, 1834) | Abundant across Vice-County | Frequent across Vice-County | 3 records, most recent 2011 |
| <i>Sciaphilus asperatus</i> (Bonsdorff, 1785) | 4 records, most recent 2016 | 5 records, most recent 2018 | 3 records, most recent 2011 |
| <i>Andrion regensteineuse</i> (Herbst, 1797) | Abundant across Vice-County | Abundant across Vice-County | Single record 2016 |
| <i>Charagmus griseus</i> (Fabricius, 1775) | 4 records, most recent 2012 | Frequent in sandy areas | Single record 1950 |
| <i>Charagmus gressorius</i> (Fabricius, 1792) | | Single record 2014 | |
| <i>Coelositona cambricus</i> (Stephens, 1831) | 8 records, most recent 2016 | Abundant across Vice-County | |
| <i>Sitona cylindricollis</i> Fähræus, 1840 | 3 records, most recent 2017 | 3 records, most recent 2013 | Single record 2014 |
| <i>Sitona hispidulus</i> (Fabricius, 1777) | Frequent across Vice-County | Frequent across Vice-County | Single record 2014 |
| <i>Sitona humeralis</i> Stephens, 1831 | 4 records, most recent 2019 | Single record 2006 | Single record 2014 |
| <i>Sitona lateralis</i> Gyllenhal, 1834 | | Single record 2000 | |
| <i>Sitona lineatus</i> (Linnaeus, 1758) | Abundant across Vice-County | Abundant across Vice-County | 4 records, most recent 2017 |
| <i>Sitona lineellus</i> (Bonsdorff, 1785) | | 6 records, most recent 2017 | |
| <i>Sitona macularius</i> (Marsham, 1802) | Single record 1980 | | |
| <i>Sitona macularius</i> (Marsham, 1802) | | 2 records, 1960 & 1966 | |
| <i>Sitona obsoletus</i> (Gmelin, 1790) | Abundant across Vice-County | Abundant across Vice-County | Single record 2018 |
| <i>Sitona puncticollis</i> Stephens, 1831 | 2 records, 1921 & 1979 | Single record 2011 | |
| <i>Sitona striatellus</i> Gyllenhal, 1834 | 5 records, most recent 2015 | 7 records, most recent 2016 | |
| <i>Sitona sulcifrons</i> (Thunberg, 1798) | 6 records, most recent 2017 | 4 records, most recent 2000 | |
| <i>Sitona suturalis</i> Stephens, 1831 | Frequent across Vice-County | Frequent across Vice-County | Single record 2014 |
| <i>Caenopsis waltoni</i> (Boheman, 1842) | 2 records, 1908 & 1979 | | |
| <i>Cathormiocerus aristatus</i> (Gyllenhal, 1827) | Single record 1993 | | |
| <i>Cathormiocerus spinosus</i> (Goeze, 1777) | | 2 records, 1966 & 1987 | |

| | Preferred Species Name | VC58 Cheshire | VC59 South Lancashire | VC60 West Lancashire |
|--|---|-------------------------------------|------------------------------|----------------------|
| | <i>Romualdius angustisetulus</i> (Hansen, 1915) | Single record 2000 | | Single record 1999 |
| | <i>Tropiphorus elevatus</i> (Herbst, 1795) | | | Single record 1950 |
| | <i>Tropiphorus terricola</i> (Newman, 1838) | 6 records, most recent 2016 | Single record pre-1900 | |
| | | | | |
| | Hyperinae | | | |
| | <i>Hypera arator</i> (Linnaeus, 1758) | Single record 1960 & 5 pre-1900 | | |
| | <i>Hypera melancholica</i> (Fabricius, 1792) | 2 records, 1962 & 1963 | | |
| | <i>Hypera miles</i> (Paykull, 1792) | Single record 2016 | | Single record 2016 |
| | <i>Hypera nigrirostris</i> (Fabricius, 1775) | 5 records, most recent 2015 | Frequent across Vice-County | Single record 2016 |
| | <i>Hypera plantaginis</i> (De Geer, 1775) | 5 records, most recent 2016 | Abundant across Vice-County | Single record 2011 |
| | <i>Hypera postica</i> (Gyllenhal, 1813) | 4 records, most recent 2017 | 2 records, 1998 & 2011 | |
| | <i>Hypera venusta</i> (Fabricius, 1781) | 3 records, 1989 and 2 pre-1900 | 3 records, most recent 1998 | |
| | <i>Brachypera dauci</i> (Olivier, 1807) | | Single record pre-1900 | |
| | <i>Brachypera zoilus</i> (Scopoli, 1763) | Single record 2016 and two pre-1900 | | |
| | <i>Hypera meles</i> (Fabricius, 1792) | Single record 2005 | Single record 2016 | |
| | <i>Hypera arundinis</i> (Paykull, 1792) | Single uncertain record 1981 | | |
| | <i>Hypera conmaculata</i> (Herbst, 1795) | Single record 2015 | Single record 2016 | |
| | <i>Hypera rumicis</i> (Linnaeus, 1758) | Frequent across Vice-County | 3 records, most recent 2012 | |
| | <i>Limobius borealis</i> (Paykull, 1792) | | | Single record 1958 |
| | | | | |
| | Lyxinae | | | |
| | <i>Cleonis pigra</i> (Scopoli, 1763) | | Frequent in sandy areas | |
| | <i>Rhinocyllus conicus</i> (Frolich, 1792) | Single record 2020 | | |
| | | | | |
| | Mesoptilinae | | | |
| | <i>Magdalis phlegmatica</i> (Herbst, 1797) | | | Single record 2011 |
| | <i>Magdalis ruficornis</i> (Linnaeus, 1758) | 2 records, 2004 & 2011 | | |
| | <i>Magdalis armigera</i> (Geoffroy in Fourcroy, 1785) | 3 records, most recent 2012 | Single record 2011 | |
| | <i>Magdalis carbonaria</i> (Linnaeus, 1758) | | 3 records, most recent 2010 | Single record 2011 |
| | <i>Magdalis cerasi</i> (Linnaeus, 1758) | 5 records, most recent 2010 | Single record 2009 | |
| | | | | |
| | Molytinae | | | |
| | <i>Leiosoma deflexum</i> (Panzer, 1795) | Abundant across Vice-County | 9 records, most recent 2018 | |
| | <i>Leiosoma oblongulum</i> Boheman, 1842 | 2 records, 1947 | | |
| | <i>Anoplus plantaris</i> (Naezen, 1794) | 6 records, most recent 2016 | 5 records, most recent 2010 | Single record 2017 |
| | <i>Anoplus roboris</i> Suffrian, 1840 | Single record 2017 | | |
| | <i>Hylobius abietis</i> (Linnaeus, 1758) | Frequent across Vice-County | Frequent across Vice-County | |
| | <i>Pissodes castaneus</i> (De Geer, 1775) | Single record 2010 | 7 records, most recent 1987 | |
| | <i>Pissodes pini</i> (Linnaeus, 1758) | | 5 records, most recent 1999 | |
| | <i>Pissodes validirostris</i> (C. R. Sahlberg, 1834) | | Single uncertain record 1953 | |
| | <i>Trachodes hispidus</i> (Linnaeus, 1758) | 3 records, most recent 2001 | | |
| | | | | |
| | Orobitidinae | | | |
| | <i>Orobitis cyanea</i> (Linnaeus, 1758) | | 8 records, most recent 2000 | |

| | Preferred Species Name | VC58 Cheshire | VC59 South Lancashire | VC60 West Lancashire |
|--|---|------------------------------|------------------------------|-----------------------------|
| | | | | |
| | Scolytinae | | | |
| | <i>Scolytus intricatus</i> (Ratzeburg, 1837) | Frequent across Vice-County | 2 records, 1990 & 2012 | |
| | <i>Scolytus laevis</i> Chapuis, 1869 | Single record 2011 | 2 records, 1981 | |
| | <i>Scolytus mali</i> (Bechstein & Scharfenberg, 1805) | Single record pre-1900 | Single record pre-1900 | |
| | <i>Scolytus multistriatus</i> (Marsham, 1802) | Single record 1984 | Single record 1993 | |
| | <i>Scolytus rugulosus</i> (P. W. J. Müller, 1818) | 2 records, 1930 & 1985 | | |
| | <i>Scolytus scolytus</i> (Fabricius, 1775) | 2 records, 1984 & 2012 | 6 records, most recent 1992 | |
| | <i>Pityophthorus pubescens</i> (Marsham, 1802) | Single record 1991 | | |
| | <i>Cryphalus asperatus</i> (Gyllenhal, 1813) | | Single record 1965 | |
| | <i>Ernoporicus caucasicus</i> (Lindemann, 1876) | 2 records, 2000 | | 4 records, 2002 |
| | <i>Trypophloeus binodulus</i> (Ratzeburg, 1837) | | Single record 1989 | |
| | <i>Dryocoetes alni</i> (Georg, 1856) | 5 records, most recent 1987 | 2 records, 1987 and pre-1900 | |
| | <i>Dryocoetes autographus</i> (Ratzeburg, 1837) | Single record 1985 | | |
| | <i>Dryocoetes villosus</i> (Fabricius, 1792) | Frequent across Vice-County | Frequent across Vice-County | |
| | <i>Ips acuminatus</i> (Gyllenhal, 1827) | | Single record pre-1900 | |
| | <i>Ips cembrae</i> (Heer, 1836) | | | Single record 1991 |
| | <i>Ips sexdentatus</i> (Boerner, 1767) | Single record pre-1900 | Single record pre-1900 | |
| | <i>Orthotomicus erosus</i> (Wollaston, 1857) | | Single record 1993 | |
| | <i>Orthotomicus laricis</i> (Fabricius, 1792) | Single record pre-1900 | | |
| | <i>Orthotomicus suturalis</i> (Gyllenhal, 1827) | 2 records, 2018 | | |
| | <i>Pityogenes bidentatus</i> (Herbst, 1784) | 2 records, 1965 & 1991 | | |
| | <i>Pityogenes chalcographus</i> (Linnaeus, 1760) | Single record 1900 | | |
| | <i>Xyleborinus saxesenii</i> (Ratzeburg, 1837) | 6 records, most recent 1992 | 4 records, most recent 1992 | |
| | <i>Anisandrus dispar</i> (Fabricius, 1792) | | Single record 1987 | |
| | <i>Trypodendron domesticum</i> (Linnaeus, 1758) | Frequent across Vice-County | Frequent across Vice-County | 6 records, most recent 2017 |
| | <i>Trypodendron lineatum</i> (Olivier, 1795) | 2 records, 1986 & 2011 | | |
| | <i>Trypodendron signatum</i> (Fabricius, 1792) | Single record 1986 | | |
| | <i>Hylesinus crenatus</i> (Fabricius, 1787) | 2 records, 1984 & 1991 | 3 records, most recent 1990 | |
| | <i>Hylesinus taranio</i> (Danthoine in Bernard, 1788) | 2 records, 1988 | | |
| | <i>Hylesinus varius</i> (Fabricius, 1775) | Single record 2013 | Single record 2014 | 3 records 2011 |
| | <i>Kissophagus vicinus</i> (Comolli, 1837) | 2 records, 2000 and pre-1900 | | |
| | <i>Pteleobius vittatus</i> (Fabricius, 1792) | Single record 1979 | | |
| | <i>Hylastes angustatus</i> (Herbst, 1794) | Single uncertain record 2015 | | |
| | <i>Hylastes ater</i> (Paykull, 1800) | 9 records, most recent 2018 | 4 records, most recent 1986 | Single record 2006 |
| | <i>Hylastes brunneus</i> (Erichson, 1836) | Single record 1950 | | |
| | <i>Hylastes cucicularius</i> Erichson, 1836 | Single record 1991 | | |
| | <i>Hylastes opacus</i> Erichson, 1836 | Single record 1974 | Single record 1987 | |
| | <i>Hylurgops palliatus</i> (Gyllenhal, 1813) | 3 records, most recent 1991 | 2 records, 1965 & 2002 | |
| | <i>Tomicus piniperda</i> (Linnaeus, 1758) | 4 records, most recent 1991 | 8 records, most recent 1996 | 2 records, 1992 & 1995 |
| | <i>Phloeotribus rhododactylus</i> (Marsham, 1802) | | 2 records, 1998 | |